

Monoferric® *ferric derisomaltose*

Monoferric® is an intravenous (IV) iron treatment for iron deficiency or iron deficiency anemia. It is used when oral iron is not effective, not tolerated, or when rapid iron replacement is needed. It can be used in non-pregnant patients and during pregnancy (second and third trimester) when recommended by your healthcare provider.

WHY YOU MIGHT NEED IV IRON

- You cannot tolerate oral iron
- You have moderate to severe iron deficiency anemia
- You need rapid iron replacement
- You have ongoing blood loss or poor iron absorption
- You are pregnant and need faster correction of anemia

USE DURING PREGNANCY

- Oral iron is not effective or causes side effects
- Anemia is moderate to severe
- Iron replacement is needed later in pregnancy

HOW IT IS GIVEN

- Given through a vein (IV infusion)
- Usually given as a single full-dose infusion
- Infusion typically takes 20–60 minutes, depending on the dose
- You will be monitored during and briefly after the infusion

AFTER YOUR INFUSION

- You can usually return to normal activities the same day
- Symptoms may improve over days to weeks
- Bloodwork is repeated 4–8 weeks after treatment to check your response

COMMON SIDE EFFECTS

Most are mild and temporary.

- Headache
- Nausea
- Flushing
- Dizziness
- Muscle or joint aches

LESS COMMON BUT IMPORTANT

Tell your care team if these occur.

- Allergic reactions
- Low blood pressure during infusion
- Skin staining if iron leaks outside the vein

SEEK URGENT MEDICAL ATTENTION IF YOU HAVE:

- Difficulty breathing
- Chest tightness
- Swelling of the face or throat
- Severe rash

If you are pregnant and have a reaction to IV iron, you will be sent to the emergency department for assessment.

WHEN TO CONTACT YOUR CARE TEAM

Contact Ferra (or your healthcare provider) if you experience:

- Fever
- Increasing pain or swelling at the IV site
- Persistent rash or shortness of breath
- Any concerning symptoms after treatment